

**KILLS THE
ROOT**

**MONEY BACK
GUARANTEE***

**Kills Listed Weeds
NOT The Lawn***

Spectracide®

**WEED STOP®
FOR LAWNS**

**Plus
CRABGRASS KILLER**

CONCENTRATE

Herbicida concentrado para céspedes y contra mano de cangrejo

**Kills 470+ Weed
Types As Listed**

**Visible Results
In 5 Hours**

Kills Yellow Nutsedge

For Best Results, Apply When Temperature Is Between 45°F - 90°F

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt.....	3.74%
Quinclorac.....	1.79%
Dicamba, dimethylamine salt.....	0.43%
Sulfentrazone.....	0.22%
OTHER INGREDIENTS.....	93.82%
TOTAL	100.00%

(For terms of guarantee, see back panel.)
*When used as directed

THIS PRODUCT CONTAINS: 0.264 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 3.11%; 0.152 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 1.79%; 0.030 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.36%; 0.019 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.22%. Isomer Specific by AOAC Method.

**KEEP OUT OF REACH OF CHILDREN
CAUTION** See back panel for additional
Precautionary Statements.

**MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN** Consulte el folleto en la parte posterior para
conocer las declaraciones preventivas adicionales.



**CONNECT
TO HOSE**

**Net Contents 32 fl oz
(1 qt / 946 mL) 17-16186**

DECLARACIONES PREVENTIVAS

Riesgos para los seres humanos y los animales domésticos

PRECAUCIÓN. Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Nocivo si se ingiere. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines, zapatos y guantes de goma/hule.

Después de manipularlo, lávese muy bien con agua y jabón. Después de utilizar este producto, enjuague los guantes antes de quitárselos, quítese la ropa y lávela por separado antes de volver a usarla y rápidamente lávese muy bien las manos y la piel que estuvo expuesta, con agua y jabón. Quítese la ropa mojada y báñese en regadera tan pronto como sea posible.

Primeros auxilios

Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15 a 20 minutos. • Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
Si se ingiere:	• Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito a menos que así lo indique un centro de control de envenenamientos o el médico. • No administre nada a una persona que haya perdido el conocimiento.

Quando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-800-917-5438 para obtener información acerca del tratamiento médico de urgencia.

Riesgos para el medio ambiente

Este pesticida es tóxico para los peces y los invertebrados acuáticos. El arrastre y el escurrimiento pueden resultar peligrosos para los organismos acuáticos de las zonas cercanas al sitio de aplicación. Para proteger el medio ambiente, no permita que el pesticida entre o se escurra por drenajes pluviales, zanjas de drenaje, cunetas ni aguas superficiales. Aplicar este producto cuando no haya viento y no se pronostique lluvia en las siguientes 24 horas ayudará a asegurar que ni el viento ni la lluvia arrastren o quiten el pesticida de la zona de tratamiento. Enjuagar el equipo de aplicación encima de la zona tratada ayudará a evitar el escurrimiento a masas de agua o sistemas de drenaje.

Riesgos físicos y químicos

Combustible. No lo use ni almacene cerca de una fuente de calor ni del fuego.

NOTA: Hasta el grado que sea compatible con la legislación pertinente, el comprador asume toda la responsabilidad para seguridad y uso que no estén de acuerdo con las instrucciones.

¿Preguntas o comentarios? Llame al 1-800-917-5438.

Spectracide

WEED STOP FOR LAWNS PLUS CRABGRASS KILLER CONCENTRATE

Herbicida concentrado para céspedes y contra mano de cangrejo

PEEL HERE • RESEALABLE LABEL
ABRA AQUÍ • ETIQUETA RESELLABLE

WHERE TO USE



Lawns — Treats Bluegrass,
Fescue, Zoysia**, Bermuda***,
Perennial Ryegrass

WHEN TO USE

45°F
90°F

For best results, apply between 45°F–90°F.
Above 45°F, the better the weeds will be
able to absorb the product.
Lawn damage could happen above 90°F.

KILLS ALL TYPES OF LISTED BROADLEAF & GRASSY WEEDS

Including Crabgrass, Dandelion, Chickweed, Clover & Yellow Nutsedge



Kills Listed Weeds Including
Crabgrass — Not the Lawn*
*When used as directed



Kills 470+ Weed
Types as Listed



Visible Results
in 5 Hours



Kills by Contact



Kills Weed Roots!



Treats up to 5,000 sq ft

**Do not apply to zoysiagrass just emerging from dormancy.

***Application to Bermudagrass may cause temporary yellowing or discoloration, but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 90°F.

WHAT TO EXPECT



PRE-APPLICATION

- **PICTURED WEED:**
Plantain
- **APPLICATION RATE:**
7 fl oz/gallon of water



5 HOURS

- **EXPECTED VISUAL SYMPTOMS:** Leaves yellowing/browning, then curling, twisting, wilting & shriveling
- *Visual symptoms and time to kill will depend on weed type, temperature and application rate



24 HOURS



96 HOURS

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

QUESTIONS & COMMENTS: Call 1-800-917-5438 or visit our website at www.spectracide.com

NOTA: Hasta el grado que sea compatible con la legislación pertinente, el comprador asume toda la responsabilidad para seguridad y uso que no estén de acuerdo con las instrucciones.

¿Preguntas y comentarios? Llame al 1-800-917-5438.

***GUARANTEED RESULTS ON YOUR MONEY BACK**

If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

Distributed by Spectrum Group, Division of United Industries Corporation

PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 9688-268-8845

EPA Est. No. 9688-MO-1 (D) 58906-MO-1 (A-E)

Circled letter is first letter of lot number.

17-15281 © 2018 UIC

PROOF OF PURCHASE

FPO UPC

0 71121 95703 0

STOP READ ENTIRE LABEL BEFORE USE.

ALTO. LEA TODA LA ETIQUETA ANTES DE USAR.

CONTROLA LA MAYORÍA DE LAS MALEZAS COMUNES, INCLUYENDO:

Alder	Big leaf maple	Broomweed	Common cocklebur	Cudweed (Purple cudweed)	Field horsetail (Common horsetail,
Alfalfa (Lucerne)	Bigroot morning glory	Buckwheat	(Rough cocklebur,		Mare's tail)
Alpine aster	Bigtooth aspen	Buttobush	Large cocklebur,	Cupid shaving brush	
Alsike clover	(American aspen,	Bull mallow (French mallow)	Common dandelion	Curly dock (Curled dock, Yellow dock,	Field mouse-ear (Field chickweed)
Alyce clover	White poplar)	Bull nettle	Common groundsel	Sour dock	Field oxeye-daisy
American black elderberry	Bi-lobed speedwell (Twolobe speedwell)	Bull thistle	Common hawkweed (Yellow hawkweed)	Curly indigo	(Creeping oxeye, Oxeye daisy)
American elm (White elm)	Bindweed (Wild morning glory)	Bur oak	Common hawthorn	(Sensitive jointweed)	Field pansy
American hazel	Bird vetch	Burclover	Common honeysuckle		
American persimmon (Common persimmon)	Bitter sneezeweed Bitterweed	Burning nettle (Stinging nettle)	(Woodbine)	Curlycup gumweed	Field pennycress (French weed)
	Black bent (Redtop)	Bushy aster	Common iceplant	Cutleaf evening primrose	Field pepperweed (Field pepperwort)
American speedwell (American brooklime)	Black birch (River birch, Water birch)	Bushy buttonweed	Common knotgrass (Knottedweed)	Daisy fleabane (annual)	Field violet
American yellowrocket	Black cherry	Buttonweed (Rough buttonweed, Poorjoe)	Common lespeeza	Dayflower	Flatwoods plum (Sloe plum)
Amur honeysuckle	Black cottonwood	California knotweed	Common lupine	Dead nettle	
Annual knawel (German knotweed)	Black medick	California rose	Common mallow	Desert wild grape	Flax weed
	Black mustard	California wild grape	(Alkali mallow,	Dewberry	Floodman thistle (prairie)
	Black willow	Canada thistle	Buttonweed,	(Garden dewberry)	
Annual nettle (Dwarf nettle)	Blackberry Blackberry elder	Carolina geranium (wild)	Cheeseweed, Dwarf mallow)	Dichondra (Carolina ponysoot)	Florida betony Florida prickly blackberry
Annual rape (Wild rape)	Black-eyed Susan	Carpeweed	Common mullein	Distaff thistle	
Annual sowthistle (common sowthistle, Milk sowthistle)	Blackseed plantain	Catchweed	Common primrose (English primrose)	Dogbane	Florida pusley
	Blessed thistle	Catchweed bedstraw	(English primrose)	Dogfennel	Florida yellow woodsoorrel
	Bloodflower	Catnip	Common purslane	Dollarweed	
	(Tropical milkweed,	Catsear (Catsear dandelion, False dandelion, Spotted weed)	Common ragweed Common salsify (Goatsbeard)	(Manyflower marshpennywort)	French horn Garden spurge Garden
Annual trampweed	Mexican butterfly weed)	Common catsear)	Common snowberry	Doveweed	star-of-Bethlehem (Grass lady)
Annual wild geranium (Spotted geranium, Wood geranium, Cranesbill)	Blue ash Blue elderberry Blue lettuce Blue mustard	Common speedwell Common sunflower Common tarweed	Common speedwell Common sunflower Common tarweed	Elk thistle English daisy Eucalyptus	Garlic mustard Giant ironweed Giant ragweed Gooseberry
Annual yellow sweet clover	Blue toadflax Blue vervain Blue woodsorrel	Common thistle (Bull thistle)	Common yellow woodsorrel (Yellow woodsorrel, Common yellow oxalis)	(Blue-gum, Tasmanian blue-gum, Southern blue-gum)	Gray chickweed (Gray mouse-ear chickweed)
Arkansas rose	Blue woodsoorrel	Chicory	Common yellow woodsorrel,	Evergreen blackberry	
Artichoke thistle	Blunt-leaved milkweed	Chinese yarrow (Siberian yarrow)	Common yellow oxalis)	Fall dandelion	Gray thistle
Asiatic pennywort	(Clasping milkweed)	Chinkapin oak	Coral bead	False flax	Great burdock
Asiatic witchweed	Box elder	Choke cherry	Common chamomile	False sunflower	Great ironweed
Atlantic white cedar (Southern white cedar)	Bracted plantain Brambles	Coffeebean	Corn chamomile	Fennel	Great yellow woodsoorrel
	Brass buttons	Coffeeweed	Corn speedwell	Field bindweed	
Austrian fieldcress	Brazil pusley	(Colorado river-hemp)	Corn spurry (Morning glory, Creeping Jenny)	(Morning glory, Creeping Jenny)	Ground ivy (Creeping Charlie, Creeping Jenny)
Ball mustard	Brazilian pepper	Common burdock	Coyote brush	Field burnweed	
Bedstraw	Bristly mallow	Common cinquefoil	Creeping buttercup	(Lawn burnweed,	Groundsel
Beggarweed (Creeping beggarweed)	Bristly oxotunge Broadleaf knotweed Broad-leaved dock	(Oldfield cinquefoil)	Creeping woodsorrel Crimson clover	Common soliva, Spurweed)	Hackberry Hairy beggarticks

Hairy bittercress	Large flower pusley	Northern pin oak	Prickly sida	Sagebrush	Southern crabgrass
Hairy buttercup	Late goldenrod	Oakleaf fleabane	Prostrate knotweed	Salmonberry	Southern wild rose
Hairy fleabane	(Canada goldenrod)	Old world diamond	Prostrate pigweed	Salicedar	(Virginia rose,
Hairy gailsnoga	Leaky spurge	flower (Oldenlandia)	Prostrate spurge	Sassafras	Common wild rose
Hairy vetch	Leathy thistle	Flattop	Prostrate verbain	(White sassafras,	Spanish needles
(Fodder vetch,	Lilac	mille grains)	Prostrate vervain	Red sassafras,	Spiny amaranth
Winter vetch)	Little bittercress	Orange hawkweed	Puncture vine	Silly sassafras)	Spiny cocklebur
Hairy-pitted stork's bill	Little evening primrose	Oriental cocklebur	(Goat's-head)	Scarlet pimpernel	Spiny sowthistle
Healall	(Small sundrops)	Pale smartweed	Purple amaranth	Scotch broom	Spotted knapweed
Heartleaf drymary	Little mallow (Least	Parsley-piert	Purple aster	Scotch thistle (Cotton	Spotted loonweed
Heartleaf nettle	mallow)	(Slender	Purple cudweed	thistle)	(Freckled milkvetch)
Heath aster	Locust	parsley-piert)	(Annual purple	Scouler's willow	Spotted spurge
(White heath aster)	London rocket	Parsnip (Wild parsnip)	cudweed, Perennial	Shepherd's purse	Sprawling horseweed
Hedge mustard	Mackenzie willow	Pearlwort	purple cudweed)	Shiny cudweed	St. Johnswort
Hemp	Madrone	(Bird's-eye	Purple deadnettle	Shinny downfox tail	Sticky chickweed
Hemp sesbania	Marcela	pearlwort)	Purple milkweed	(Orange foxtail)	(Stick mouse-ear
Henbit	Marshelder	Pennsylvania	Purple milkvetch	Shortstalk stinkweed	chickweed)
Himalayan blackberry	Matchweed	cinquefoil (Prairie	(Purple loco,	Showy evening	Stiff goldenrod
Hoary cress	Mayweed	cinquefoil)	Field milkvetch)	primrose	(Stiff-leaved
Hoary plantain	(Stinking chamomile)	Pennsylvania	Purple nutsedge	Showy goldenrod	goldenrod)
Hoary vervain	Meadow Hawkweed	smartweed	(Purple nutsgrass)	Showy tick-trefoil	Strawberry clover
Hop clover	Meadow thistle	Pepperweed	Purslane speedwell	(Canadian tick-trefoil,	Swamp smartweed
Horsenettle	Mesquite (Honey	(Perennial	Quaking aspen	Canada tickclover)	Sweet goldenrod
Horseradish	mesquite)	pepperweed)	Rabbit foot clover	Silver ponysoot	Sweet gum
Horseweed/marestail	Mexican morning glory	Pigweed	Raspberry	(Silver nickel vine)	(American
Illinois woodcock	Mexicanweed	Pin oak	(Wild black	Skunkbush sumac	sweetgum)
Indian mock strawberry	(Mexican fireweed,	(Swamp Spanish	raspberry)	Slender amaranth	Swinecress
Indian mustard	Kochia)	oak)	Red alder	Slender bush clover	Tall beggaricks
Innocence (Blue-eyed	Mimosa	Pineappleweed	Red clover	(Slender lespedeza)	Tall milkweed
Mary)	Mock strawberry	Pineywoods bedstraw	Red maple	Slender plantain	(Poke milkweed)
Italian thistle	Mojave stinkweed	(Wood bedstraw)	(Water maple, Soft	Slender speedwell	Tall morning glory
lyleaf morning glory	Moneyswort	Plains coreopsis	maple)	Slender stinkweed	(Common
lyleaf speedwell	Mountain bush	(tickseed)	Red morning glory	Slender-flower thistle	morning glory)
Japanese honeysuckle	honeysuckle	Plantain	Red sorrel (Sheep's	Smallowflower buttercup	Tall nettle
Jewel milkweed	Mountain woodsorrel	(Broadleaf plantain,	sorrel)	Smallowflower gailsnoga	Tall thistle
(Pallid milkweed)	(Wood shamrock)	Greater plantain,	Redroot	Small-flowered	Tall vervain
Jeweledweed	Mouse-ear chickweed	Common plantain)	Redroot pigweed	winter-cress	Tanokak
Jimsonweed, devil's	Mouse-ear hawkweed	Platte thistle	Red-seeded dandelion	Smooth bedstraw	(Tanbark-oak)
snare	Mugwort	Plumless thistle	Redstem filaree	(Crosswort)	Tansy mustard
Johnny-jump violet	Multiflora rose	Poison hemlock	Redweed	Smooth beggaricks	Tansy ragwort
Kudzu	Musk thistle	Poison ivy	Redwood sorrel	Smooth cat's ear	Tanweed (Water
Ladythumb	Narrow leaved willow	Poison oak	(Oregon oxalis)	Smooth chaff-flower	knotted, Water
smartweed	Narrowleaf cudweed	Poison smartweed	Rough cinquefoil	Smooth crabgrass	smartweed)
Lamb's quarters	Narrowleaf plantain	Pokeweed	Rough fleabane	Smooth dock	Teaweed
(Goosefoot, Pigweed)	(English plantain,	Port Orford cedar	Rough horsetail	Smooth pigweed	Texas filaree
Lambsquarters	buckhorn)	(Lawson cypress)	(Scouringrush	Sneezeweed	Thimbleberry
Lanceleaf ragweed	Nightshade (Silverleaf)	Povertyweed	horsetail)	Snow speedwell	Three flower
Lantana	Northern bedstraw	Prairie sunflower	Russian pigweed	Snow	beggarweed
Large crabgrass	Northern bush	Prickly lettuce	Russian thistle	Southern bush	Thymeleaf
(Hairy crabgrass)	honeysuckle	(Compass plant)	Sage	honeysuckle	speedwell

Toadflax (Common toadflax, Yellow toadflax)	Vetch (Common vetch, Garden vetch)	White ash (American ash)	Wild honeysuckle (Tartarian honeysuckle)	Winter speedwell (Persian speedwell)	Yellow birch (Golden birch)
Tooth-leaved croton (Tropic croton, Sand croton)	Vine maple	White clover ("Dutch clover, Honeysuckle clover, White trefoil, Purplewort)	Wild lettuce	Wood nettle	Yellow nutsedge
Trailing blackberry	Violet woodsorrel	White mustard	Wild marigold (Hogwort, Goatweed)	Woolly croton	Yellow rocket (Winter cress, Bitter wintercress,
Trailing crownvetch	Virginia creeper	White starwort	(Roundleaf marigold, Southern marigold)	Woolly morning glory	Wild mustard)
Tree tobacco (Peppergrass)	Virginia pepperweed	White prairie aster	Wild morning glory (hedge bindweed)	Woollyleaf bursage	Yellow starthistle
Trumpet creeper (Trumpet vine, Cow itch vine, Hummingbird vine)	Virginia winged rockcress	White sweet clover (White melilot)	Wild plum (American plum)	Woolly burdock (Downy burdock)	Yellow sundrops (Shrubby evening primrose)
Tufted evening primrose (Fragrant evening primrose)	Wandering cudweed	Whitestem filaree	Wild radish	Wooly locweed	Yellow sweet clover (Yellow melilot, Ribbed melilot, Common melilot)
Tufted knotweed	Water pennycort (Floating marshpennywort)	Whitetop	Wild rose (Prickly wild rose)	Wormseed	Yellow thistle (horrible yellowflower pepperweed
Tumble mustard	Wayleaf thistle (Gray thistle)	Wild blackberry (Thornless blackberry)	Wild strawberry	Wormseed mustard	Yellowspine thistle
Tumble pigweed	Western clematis	Wild buckwheat	Wild sweet potato	Wreath goldenrod (Woodland goldenrod)	*annual or perennial
Velvetleaf	Western ragweed	Wild carrot	Wild vetch	Wrinkleleaf goldenrod (Roughstemmed goldenrod)	
Venice mallow	Western salsify (Goatsbeard)	Wild cherry	Wild violet		
		Wild four-o'clock	Willow-leaved poplar (Narrowleaf cottonwood)	Yarrow (Common yarrow)	
		Wild garlic (Wild onion)			

STOP. READ ENTIRE LABEL BEFORE USING.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. *For outdoor residential use only.*

General Precautions and Restrictions

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried. Do not exceed more than 0.375 lb Sulfentrazone/A per year. Do not treat when air temperatures exceed 90°F, as damage to grass may occur. Application at rates above those specified on this label may cause injury to turfgrass. Do not apply this product to food crops (gardens and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other non-target plants. Do not spray exposed roots of ornamentals and trees.

Do not use Spectracide® Weed Stop® For Lawns Plus Crabgrass Killer Concentrate on lawns containing bentgrass, St. Augustinegrass, bahiagrass, centipedegrass or carpetgrass. Also, do not use on dichondra, or lawns with desirable clovers or legumes. Do not use on vegetable gardens, fruits or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs).

WHERE TO USE

This product is intended for use by homeowners on residential lawns. Only apply this product to the following species of turfgrass:

Cool season turfgrass: Kentucky bluegrass; perennial ryegrass; fescue spp., including tall, red and fine leaf fescues.

Warm season turfgrass: zoysiagrass^{**}, Bermudagrass^{***}, buffalograss

**Do not apply to zoysiagrass just emerging from dormancy, as it may result in lawn damage.

***Application to Bermudagrass may cause temporary yellowing or discoloration, but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 90°F.

WHEN TO USE#

Apply when daytime temperatures are between 45°F and 90°F

SPRING: Spray when broadleaf weeds are young and actively growing with adequate soil moisture for best results

FALL: Spray when broadleaf weeds are young and actively growing with adequate soil moisture for best results. Fall applications will control weeds that may otherwise go dormant through the winter and resprout the following spring. For heavy grass weed pressure, apply a second application at the same rate, 14 days after the first application.

CRABGRASS (LARGE AND SMOOTH): Spring and early summer: For the best results, apply this product from crabgrass emergence until the six-leaf stage (two tillers) or before the crabgrass is 3 to 4 inches tall. This period begins the first time that you can see crabgrass in your lawn. Use two applications at the same rate with a 14-day interval for dense populations.

FOXTAIL (GREEN, GIANT AND YELLOW):

Spring and early summer: For the best results, apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches tall. Use two applications at the same rate with a 14-day interval for dense populations.

YELLOW NUTSEDGE:

Early summer: Two applications of this product are required when the yellow nutsedge plants are 3 to 6 inches tall. Apply the second application at the same rate 14 days later.

BROADLEAF WEEDS:

Spring: Apply when dandelion, plantain, buttercup, speedwell, knotweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

Fall: Apply when henbit, chickweed, white clover, ground ivy, lawn burweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

**Visual symptoms and time to kill will depend on weed type, temperature and application rate.*

HOW MUCH TO USE

See the following table for calculating the correct amount of product to use for the area to be treated. The maximum application rate is 6.5 fl oz of product per 1,000 sq ft per application (0.6 lb 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to two per year. The maximum seasonal rate is 13 fl oz of product per 1,000 sq ft (1.2 lb 2,4-D acid equivalent per acre), excluding spot treatments.

MEASUREMENT CALCULATIONS: Total lawn area in square feet = length x width.

DIRECTIONS FOR HOSE-END SPRAYER

1. Shake container well before using.
2. Connect a garden hose to the sprayer nozzle. Make sure switch is fully forward in the "OFF" position.
3. Turn on water at faucet. Extend hose to the farthest area to be treated and work back toward the faucet so you don't come in contact with the treated area.
4. Remove safety tab from sprayer's right side by pulling straight out. Keep for future reuse.
5. To BEGIN spraying, point nozzle toward treatment site and pull switch backwards with thumb toward hose connection. Water will automatically mix with the product.
6. Spray until wet to control weeds. Walk at a steady pace, working back toward the faucet while spraying, using an even sweeping motion, slightly overlapping treated areas. One quart will treat 5,000 sq ft of Southern or Northern grasses.
7. To STOP spraying, push switch forward with thumb toward nozzle. Turn water off at faucet. To relieve pressure before removing nozzle from hose, pull switch backwards with thumb toward hose connection until water stops spraying.
8. To STORE unused product, make sure switch is fully forward in the "OFF" position.
9. Reinstall safety tab into sprayer's right side. With sprayer in the full "OFF" position, push tab into position.
10. Place in cool area away from heat, sunlight or open flame.

Turfgrass	Total treated area for 1 qt (32 fl oz) sprayer	Amount of herbicide applied per 1,000 sq ft
Kentucky bluegrass, fescue spp., perennial ryegrass, zoysiagrass**, Bermudagrass*** and buffalograss	5,000 sq ft	6.4 fl oz

**Do not apply to zoysiagrass just emerging from dormancy, as it may result in lawn damage.

***Application to Bermudagrass may cause temporary yellowing or discoloration, but full recovery can be expected. Do not apply to Bermudagrass when daytime temperatures exceed 90°F.

USE TIPS

- Adequate moisture and mild environmental conditions enhance the performance of this product. Applications should be made to actively growing weeds. Apply when daytime temperatures are between 40°F and 90°F. Avoid applications of this product when turfgrasses are under stress since injury may result.
- **For newly seeded areas:** Do not apply within four weeks after seedling emergence of Kentucky bluegrass, fine fescue blends and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately six weeks after seedling emergence) or after the third mowing.
- **For newly sodded, sprigged or plugged areas:** The application of this product to newly sodded, sprigged or plugged grasses should be delayed until three to four weeks after the sodding, sprigging or plugging operations.
- **Seeding:** Turf species listed on this label can be seeded into the treated areas at four weeks after the application of this product.
- **Irrigation:** Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application. If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended.
- **Mowing:** Delay mowing two days before and until two days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury. Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals or trees in vegetable gardens.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry area away from heat or open flame. **Pesticide Disposal and Container Handling:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

Wash thoroughly with soap and water after handling. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If swallowed:	• Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a Poison Control Center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-800-917-5438 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

Physical and Chemical Hazards

Combustible. Do not use or store near heat or open flame.

ALTO. LEA TODA LA ETIQUETA ANTES DE USAR.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto de una manera que no sea la indicada en la etiqueta. Solo para uso residencial en espacios exteriores.

Precauciones y restricciones generales

No aplique este producto de modo que entre en contacto con cualquier persona o mascota, ya sea directamente o por arrastre. Mantenga a las personas y las mascotas fuera de la zona durante la aplicación. No permita que las personas ni las mascotas entren en la zona tratada hasta que los rocíos se hayan secado. No exceda 0.375 lb de sulfentrazone/A por año. No trate cuando la temperatura del aire sea superior a 90°F porque se puede dañar el césped. La aplicación de dosis superiores a las que se especifican en esta etiqueta puede causar daño al césped. No aplique este producto a cultivos de alimentos (jardines y hortalizas), cultivos de forraje ni plantas ornamentales, incluyendo flores, árboles, arbustos, setos, plantas ornamentales leñosas, cobertura de suelo establecida en plantaciones ornamentales y otras plantas que no sean el objetivo. No rocíe las raíces expuestas de árboles y plantas ornamentales.

No use el Herbicida concentrado para céspedes y contra mano de cangrejo Spectracide® Weed Stop® en céspedes que contengan agrostis, pasto San Agustín, bahiagrass, ciempiés o césped alfombra. Además, no lo use en dichondra ni en céspedes con tréboles o legumbres deseables. No use en huertos, frutas o plantas ornamentales (flores, árboles, cobertura de suelos, setos, jardines/lechos de flores y arbustos).

DÓNDE USAR

Este producto está diseñado para uso de los propietarios en céspedes residenciales. Aplique este producto únicamente a las siguientes especies de céspedes:

Césped de clima frío: pasto azul de Kentucky; ryegrass perenne; especies de festuca, incluyendo alta, roja y de hoja fina.

Césped de clima cálido: zoysia**, Bermuda*** y buffalograss

**No aplique al césped zoysia recién saliendo de inactividad, ya que puede causar daño al césped.

***La aplicación al césped Bermuda puede causar amarillamiento o decoloración temporal, pero se puede esperar una recuperación total. No aplique al césped Bermuda cuando las temperaturas diurnas sean superiores a los 90°F.

CUÁNDO USAR‡

Aplique cuando las temperaturas diurnas se encuentren entre 45°F y 90°F.

PRIMAVERA: Para obtener mejores resultados, rocíe cuando las malezas de hoja ancha son jóvenes y están en crecimiento activo con la humedad de suelo adecuada.

OTOÑO: Para obtener mejores resultados, rocíe cuando las malezas de hoja ancha son jóvenes y están en crecimiento activo con la humedad de suelo adecuada. Las aplicaciones en otoño controlarán las malezas que, de otra manera, podrían pasar inactivas en invierno y volver a brotar en la primavera siguiente. Si hay muchas malezas herbosas, haga una segunda aplicación con la misma

dosis, 14 días después de la primera aplicación.

MANO DE CANGREJO (GRANDE Y SUAVE): Primavera y principios del verano: Para obtener mejores resultados, aplique este producto desde la emergencia del pasto mano de cangrejo hasta la etapa de seis hojas (dos retoños) o antes de que el pasto mano e cangrejo tenga de 3 a 4 pulgadas de altura. Este período comienza la primera vez que usted observa mano de cangrejo en su césped. Haga dos aplicaciones con la misma dosis con un intervalo de 14 días para poblaciones densas.

COLA DE ZORRO (VERDE, GIGANTE Y AMARILLA):

Primavera y principios del verano: Para obtener mejores resultados, aplique este producto a la cola de zorro durante las etapas de crecimiento inicial o cuando la planta tenga de 1 a 4 pulgadas de altura. Haga dos aplicaciones con la misma dosis con un intervalo de 14 días para poblaciones densas.

CYPERUS ESCULENTUS (JUNCIA AVELLANADA):

Principios del verano: Se necesitan dos aplicaciones de este producto cuando las plantas de juncia avellanada tienen de 3 a 6 pulgadas de altura. Realice la segunda aplicación con la misma dosis 14 días después.

MALEZAS DE HOJA ANCHA:

Primavera: Para obtener mejores resultados, aplique cuando el diente de león, llantén, ranúnculo, verónica, polígono y otras malezas de hoja ancha son jóvenes y están en crecimiento activo con la humedad de suelo adecuada.

Otoño: Para obtener mejores resultados, aplique cuando la ortiga mansa, pampina, trébol blanco, hiedra terrestre, Soliva sessilis y otras malezas de hoja ancha son jóvenes y están en crecimiento activo con la humedad de suelo adecuada.

‡Las señales visuales y el tiempo que tarda la eliminación dependerá del tipo de maleza, la temperatura y la dosis de aplicación.

CUÁNTO USAR

Vea la siguiente tabla para calcular la cantidad correcta de producto que debe usar para el área a tratar.

La dosis máxima de aplicación es de 6.5 onzas líquidas de producto por cada 1,000 pies cuadrados por aplicación (0.6 lb de equivalente ácido 2,4-D por acre, por aplicación).

La cantidad máxima de aplicaciones al voleo está limitada a dos por año.

La dosis máxima por temporada es de 13 onzas líquidas de producto por cada 1,000 pies cuadrados (1.2 lb de equivalente ácido 2,4-D por acre), excluyendo los tratamientos localizados.

CÁLCULOS DE MEDICIÓN: El área total del césped en pies cuadrados = largo x ancho.

INSTRUCCIONES PARA ROCIADORES CON APLICADOR TIPO MANGUERA

1. Agite bien el envase antes de usar.
2. Conecte una manguera de jardín a la boquilla del rociador. Asegúrese de que la llave esté completamente al frente en la posición “OFF” (apagado).
3. Abra el agua del grifo. Extienda la manguera hasta la zona más alejada que se vaya a tratar y retroceda hacia el grifo de manera de no entrar en contacto con la zona tratada.
4. Quite la lengüeta de seguridad de la parte derecha del rociador jalándola hacia afuera. Guarde para su uso futuro.
5. Para COMENZAR a rociar, dirija la boquilla hacia la zona a tratar y con el dedo pulgar, jale la llave hacia atrás hacia la conexión de la manguera. El agua se mezclará automáticamente con el producto.
6. Rocíe hasta mojar para controlar las malezas. Camine a ritmo regular, volviendo hacia el grifo mientras rocía con un movimiento de barrido parejo, superponiendo ligeramente las zonas tratadas. Un cuarto de gallón trata 5,000 pies cuadrados de céspedes del sur y del norte.
7. Para DETENER el rocío, presione con el dedo pulgar la llave adelante hacia la boquilla. Cierre el agua del grifo. Para reducir la presión antes de quitar la boquilla de la manguera, con el dedo pulgar, jale la llave atrás hacia la conexión de la manguera hasta que el agua deje de salir como rocío.
8. Para GUARDAR el producto sin usar, asegúrese de que la llave esté completamente al frente en la posición “OFF” (apagado).

9. Vuelva a colocar la lengüeta de seguridad en la parte derecha del rociador. Con el rociador en la posición “OFF” (apagado) completamente, presione la lengüeta en su posición.
10. Guarde en un lugar fresco, lejos del calor, la luz solar o el fuego.

Césped	Área total tratada para rociador de 1 cuarto de galón (32 onzas líquidas)	Cantidad de herbicida aplicada por cada 1,000 pies cuadrados
Pasto azul de Kentucky, especies de festuca, ryegrass perenne, césped zoysia**, pasto Bermuda*** y buffalograss	5,000 pies cuadrados	6.4 onzas líquidas

**No aplique al césped zoysia recién saliendo de inactividad, ya que puede causar daño al césped.

***La aplicación al pasto Bermuda puede provocar que el césped se vuelva amarillo o se decolore temporalmente pero se puede esperar una recuperación completa. No lo aplique al pasto Bermuda cuando las temperaturas diurnas superen los 90°F.

CONSEJOS SOBRE EL USO

- Las condiciones ambientales templadas y de humedad adecuada mejoran el rendimiento de este producto. Las aplicaciones se deben llevar a cabo en malezas que se encuentran en la etapa de desarrollo activo. Aplique cuando las temperaturas diurnas se encuentren entre 40°F y 90°F. Evite aplicar este producto cuando los céspedes estén sometidos a estrés ya que puede ocasionar daño.
- Para zonas recién sembradas: No lo aplique en el plazo de cuatro semanas después de la germinación de las plantas de semillero de pasto azul de Kentucky, mezclas de festuca de hoja fina y ryegrass perenne. Aplique este producto a céspedes recién cultivados cuando estén bien establecidos (aproximadamente seis semanas después de la germinación de las plantas de semillero) o después del tercer corte.
- Para zonas recién cubiertas de césped, de propagación vegetativa (“sprigging”) o con “plugs”: La aplicación de este producto a pastos recién cubiertos, de propagación vegetativa (“sprigging”) o con “plugs” se debe demorar hasta tres o cuatro semanas después de operaciones de cobertura de césped, propagación vegetativa o colocación de “plugs”.
- Siembra: Las especies de césped que se indican en esta etiqueta se pueden sembrar en las zonas tratadas cuatro semanas después de la aplicación de este producto.
- Irrigación: No aplique este producto inmediatamente antes de la lluvia o el riego. Para obtener mejores resultados, no riegue ni rocíe el césped por 24 horas después de la aplicación. Si las condiciones son secas, se recomienda programar la irrigación o el riego 24 horas antes y 24 horas después de la aplicación.
- Corte: Retrase el corte dos días antes y hasta dos días después de la aplicación de este producto. Más estrés ocasionado por la altura baja del corte puede aumentar la posibilidad del daño al césped. Los recortes de los primeros tres cortes se deben dejar en la zona tratada. No use los recortes provenientes de la zona tratada como mantillo ni abono orgánicos alrededor de flores, plantas ornamentales, árboles ni en huertos.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua ni los alimentos para personas y animales por medio del almacenamiento o la eliminación. Almacenamiento del pesticida: Guarde en un lugar fresco y seco, lejos del calor y del fuego.

Eliminación del pesticida y manipulación del envase: Envase no rellenable. No vuelva a usar ni a rellenar este envase. Si está vacío: Tirelo en la basura u ofrézcalo para reciclar, si existe esa opción. Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto sin usar por ningún drenaje interno ni externo.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry area away from heat or open flame. **Pesticide Disposal and Container Handling:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

Wash thoroughly with soap and water after handling. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If swallowed:	• Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a Poison Control Center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-800-917-5438 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

Physical and Chemical Hazards

Combustible. Do not use or store near heat or open flame.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or Comments? Call 1-800-917-5438 or visit our website at www.spectracide.com

NOTA: Hasta el grado que sea compatible con la legislación pertinente, el comprador asume toda la responsabilidad para seguridad y uso que no estén de acuerdo con las instrucciones.

¿Preguntas y comentarios? Llame al 1-800-917-5438.

Distributed by Spectrum Group, Division of United Industries Corporation, PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 9688-268-8845 EPA Est. Nos. 9688-MO-1 (U), 58996-MO-1 (AE)

Circled letter is first letter of lot number. 17-15281 © 2018 UIC